## **BIBLIOGRAPHY OF WEST INDIAN MOSSES**

## William R. Buck

Buck, William R. (New York Botanical Garden, Bronx, New York 10458-5126, U.S.A.) Bibliography of West Indian mosses. Moscosoa 5: 324-339. 1989. A bibliography of references on West Indian mosses has been compiled. 287 citations are presented.

Una bibliografía sobre los muscos (Bryophyta) de las Antillas Occidentales con 287 títulos. Es publicada antes de la flora briofítica ya en preparación por el autor y William C. Steere.


Mosses have been documented from the West Indies from as early as the late 17th century. However, most work on the group has only been floristic listings or the inclusion of West Indian species in more wide-ranging monographs. Many species have not been analyzed since their original description. Indeed, the vast majority of the following references are either descriptions of new species, floristic lists or monographs that happen to cover a few West Indian species. For this reason, a monographic flora of the West Indian mosses was proposed and is underway. All species that have been described from the West Indies, as well as many from the northern Andes—the area with the closest affinities to the Antillean moss flora— are being critically studied, described and illustrated. As a preparation to the initiation of this flora, the following bibliography was compiled so that literature records could be checked. Because of increases interest in the West Indies it seemed appropriate to share the following bibliography.

## Bibliograpy

- Allen, B. H., M. R. Crosby & R. E. Magill. 1985 [1986]. A review of the genus Stenodictyon (Musci). Lindbergia 11: 149-156.
- Anderson, L. E. & V. S. Bryan. 1958. Systematics of the autoicous species of *Ditrichum* subg. *Ditrichum*. Brittonia 10: 121-137.
- Ando, H. & M. Higuchi. 1984. Caribaeohypnum, a new genus for a northern Latin American moss, Hypnum polypterum (Mitt.) Broth. Cryptog., Bryol. Lichénol. 5:7-14.
- Andrews, A. L. 1913. Sphagnaceae. N. Amer. Fl. 15:3-31.
  - . 1923. Two additions to the Bermuda flora. Bryologist 26:6.
- Armitage, E. 1932. Bryological notes on a West Indies cruise. Bryologist 35: 23-28.
- Arzeni, C. B. 1954. The Pterobryaceae of southern United States, México, Central America, and the West Indies. Amer. Midl. Naturalist 52:1-67.

Asprey, G. F. & R. G. Robbins. 1953. The vegetation of Jamaica. Ecol. Monogr. 23: 359-412, figs. 1-39, tabs. 1-6.

- Ballet, J. 1890. La Guadeloupe. Renseignements sur l'histoire, la flore, la faune, la géologie, la minéralogie, l'agriculture, le commerce, l'industrie, la législation, l'administration. 3 vols. Basse-Terre. ["Musci ou mousses" 1(1): 469–474].
- Bartram, E. B. 1928. Mosses of Jamaica. Jamaica Naturalist 1(2): 5, 6, 15–20.
- . 1931. A review of the American species of *Daltonia*. Bull. Torrey Bot. Club 58: 31–48, pls. 3, 4.
- . 1954. Briophyta. *In:* Jiménez, J. de Js., Plantas nuevas para la ciencia, nuevas para la Hispaniola y nuevas para la República Dominicana. Publ. Univ. Santo Domingo XI, 92 bis(4): 14, 15.
- . 1955. Mosses of Dominica, British West Indies. Bull. Brit. Mus. (Nat. Hist.), Bot. 2(2): 37-49.
- . 1956. Mosses of St. Vincent, British West Indies. Rev. Bryol. Lichénol. 25: 116–119.
- Bescherelle, É. 1874. Bibliographie exotique (suite). Amérique central et Antilles. Rev. Bryol. 1: 40, 41.
- . 1876. Florule bryologique des Antilles françaises. Ann. Sci. Nat. Bot. VI, 3: 175–265.
- . 1891. Revision des Fissidentacées de la Guadeloupe et de la Martinique. Rev. Bryol. 18: 49–55.
- . 1891. Musci novi guadalupenses. Rev. Bryol. 18: 75-77.
- \_\_\_\_\_. 1894. Selectio novorum muscorum. J. Bot. (Morot) 8: 59–63, 177–179, pl. I.
- . 1898. Note sur le *Philonotula papulans C.* Müll. Rev. Bryol. 25: 89, 90.
- . 1902. Note sur les mousses nouvelle recoltées à la Guadeloupe et la Martinique par le R. P. Duss. J. Bot. (Morot) 16: 6–11.
- Biebl. R. 1965. Austrocknungsresistenz tropischer Urwaldmosse auf Puerto Rico. Protoplasma 59: 277–297.
- Bizot, M. 1965. Contribution à la flora bryologique de Cuba. Bull. Mens. Soc. Linn. Lyon 34: 306–328.
- . 1973. Contribucion à la flore bryologique de Cuba (2° article). Bryo logist 76: 340–346.
- Borgensen, F. 1919. A new moss collected in Blue Mountains, Jamaica. Bot. Tidssk. 36: 279, 280 [Anoectangium incrassatum Broth., n. sp.].
- —— & C. Raunkiaer. 1918. Mosses and lichens collected in the former Danish West Indies. Danks Bot. Ark. 2(9): 1–18. [mosses pp. 3–5, det. Brotherus].
- Boswell, H. 1887. Jamaica mosses and Hepaticae. J. Bot. 25: 45-50, 118.



- 6: 565-580 [mosses p. 579].
- Wilson. 1922. Notes on plants collected by Mr. Bucher on Pico Turquino, Cuba. J. New York Bot. Gard. 23: 91–94.
- Brotherus, V. F. 1903. Musci novi Dussiani. *In:* Urban, I., Symbolae antillanae seu fundamenta florae Indiae occidentalis 3(3): 421–429.
- . 1901–1909. Bryales. Spezieller Teil. *In:* Engler, A. & K. Prantl, Die natürlichen Pflanzenfamilien 1(3): 277–1239. Engelmann, Leipzig.
- . 1924–1925. Bryales. Spezieller Teil. *In*: Engler, A. & K. Prantl, Die natürlichen Pflanzenfamilien, ed. 2, 10: 143–478; 11: 1–542.
- Browne, P. 1756. The civil and natural history of Jamaica. London [mosses pp. 82–84, pre-binomial; same in 1789 edition].
- Bucher, G. C. 1922. Narrative of an ascent of Pico Turquino, Cuba. J. New York Bot. Gard. 23:81-91 [cf. E. G. Britton, 1922; N. L. Britton & P. Wilson, 1922].
- Buck, W. R. 1980. A re-interpretation of the Fabroniaceae: additions and corrections. J. Hattori Bot. Lab. 47: 45-55.
- . 1980. Animadversions on *Pterigynandrum* with special commentary on *Forsstroemia* and *Leptopterigynandrum*. Bryologist 83: 451–465.
- . 1982. On *Meiothecium* (Sematophyllaceae). Contr. Univ. Mich. Herb. 15: 137–140.
- . 1983. A revision of the Antillean species of *Trichosteleum* (Musci: Sematophyllaceae). Moscosoa 2: 54–60.
- . 1983. A synopsis of the South American taxa of *Fabronia* (Fabronia ceae). Brittonia 35: 248–254.
- . 1983. Nomenclatural and taxonomic notes on West Indian Sematophyllaceae, Brittonia 35: 309–311.
- . 1983. New species and new combinations in the Sematophyllum subpinnatum complex (Sematophyllaceae). Brittonia 35: 327–330.
- . 1984. *Pylaisiadelpha* replaces *Brotherella* (Sematophyllaceae). Yushania 1(2): 11–13.
- \_\_\_\_\_. 1984. Taxonomic and nomenclatural notes on West Indian Hypnaceae. Brittonia 36: 178–183.
- . 1986. Wijkia (Sematophyllaceae) en the New World. Hikobia 9: 297–303.
- & H. Crum. 1978. A re-interpretation of the Fabroniaceae with notes on selected genera. J. Hattori Bot. Lab. 44: 347–369.
- & R. R. Ireland. 1985. A reclassification of the Plagiotheciaceae. Nova Hedwigia 41: 89–125.
- & W. C. Steere. 1983. Un listado preliminar de los musgos de la Espapola. Moscosoa 2: 28-53.

- Churchill, S. P. & W. R. Buck. 1982. A taxonomic investigation of Leptotheca (Rhizogoniaceae). Brittonia 34: 1-11. Crosby, M. R. 1967. Notes on the moss flora of Puerto Rico. Bryologist 70: 122, 123. \_\_\_\_\_. 1969. A revision of the tropical American moss genus Pilotrichum. Bryologist 72: 275-383. \_\_\_\_\_. 1969. Distribution patterns of West Indian mosses. Ann. Missouri Bot. Gard. 57: 145-151. \_\_\_\_. 1971. Some additional records for the moss flora of Jamaica. Caribbean J. Sci. 11(1-2); 65, 66. . 1978. New combinations in Callicosta (Musci), the correct name for Pilotrichum. Bryologist 81: 435-437. . B. H. Allen & R. E. Magill. 1985. A review of the moss genus Hypnella. Bryologist 88: 121-129. Crum, H. 1952. Pseudosymblepharis in Middle America. Bryologist 55: 137-142. \_\_\_\_. 1960. Three new species of Fissidens. Bryologist 63: 93-98. \_\_\_\_. 1965. New moss records from Haiti. Bryologist 68: 232, 233. . 1965. Refinements on the Jamaican bryoflora. Trans. Brit. Bryol. Soc. 4: 790-793. \_\_\_\_. 1972. A taxonomic account of the Erpodiaceae. Nova Hedwigia 23: 201 - 224. \_\_\_\_\_. 1980. Comments on Sphagnum henryense. Contr. Univ. Mich. Herb. 14: 53-56. \_\_\_\_\_. 1984. Notes on tropical American mosses. Bryologist 87: 203–216. & E. B. Bartram. 1958. A survey of the moss flora of Jamaica. Bull. Inst. Jamaica, Sci. Ser. 8:1-90. \_\_\_\_ & F. J. Hermann. 1972. Sphagnum magellanicum new to the West Indies. Bryologist 75: 359, 360. \_\_\_\_ & W. C. Steere, 1956. Puerto Rican Musci. New species and new combinations. Bryologist 59: 246-255. & \_\_\_\_ & \_\_\_ . 1957. The mosses of Porto Rico and the Virgin Islands. New
- Naturalist 60: 1-51.

  D'Arcy, W. G. 1971. The island of Anegada and its flora. Atoll Res. Bull. 139: 1-21 [mosses p. 17, taken from N. L. Britton, 1916].

York Acad. Sci., Sci. Surv. Porto Rico & Virgin Islands 7(4): 395–599.

& \_\_\_\_\_ . 1958. A contribution to the bryology of Haiti. Amer. Midl.

De Sloover, J. L. 1981. Octoblepharum erectifolium Mitt. ex Williams et Steereobryon subulirostrum (Schimp. ex Besch.) G. L. Smith en Guadeloupe. Cryptog., Bryol. Lichénol. 2: 471, 472.

- Dillenius, J. J. 1741. Historia muscorum. Oxford.
- Dismier, G. 1910. Revision des *Philonotis* de l'Amérique. Bull. Soc. Bot. France IV, 10: 1-37.
- Dixon, H. N. 1920. Rhaphidostegium caespitosum (Sw.) and its affinities. J. Bot. 58: 81-89.
- Duarte-Bello, P. P. 1982. Distribución de los musgos en las regiones fitogeográficas de Cuba. Acta Bot. Cubana 7: 1–20.
- . 1982. Musgos cubanos: su presencia mundial. Acta Bot. Cubana 9: 1–19.
- . 1982. Nueva especie de *Amblytropis* (Musci: Daltoniaceae) de Cuba. Acta Bot. Cubana 13: 1-5.
- Duss, R.-P. 1903. Enumeration méthodique des muscinées des Antilles françaises. II.-Mousses. Lons-Le-Saunier.
- \_\_\_\_\_. 1904. Flore cryptogamique des Antilles françaises. Lons-Le-Saunier.
- Fife, A. J. 1982. Taxnomic and nomenclatural observations on the Funariaceae. 2. Lectotypification of *Physcomitrium* subg. *Cryptopyxis* (C. Müll.) Broth. Lindbergia 8: 75, 76.
- Fleischer, M. 1922. Kritische Revision der Carl Müllerschen Laubmoosgatungen. Hedwigia 63: 209–216.
- Florschütz, P. A. 1954. Notes on American Musci. Acta Bot. Neerl. 3:94-99.

  ——. 1955. A short survey of the American species of the genus *Octoble-pharum*. Mitt. Thüring. Bot. Ges. 1: 51–58.
- . 1967. Mosses from the Netherlands Antilles. Acta Bot. Neerl. 15: 524-529, pl. I.
- Foucault, B. de 1977. Flore des bryophytes de Guadeloupe. Basse-Terre.
- Frahm, J.-P. 1975. Conspectus der mittel- und südamerikanischen *Campylopus*-Arten (Dicranaceae). Bryophyt. Biblioth. 5: 1-143, figs. 1-14.
- . 1980. Taxonomische Notizen zur Gattung *Campylopus* IX. J. Bryol. 11:213–218.
- . 1981. Taxonomische Notizen zur Gattung Campylopus X. Lindbergia 7: 27-31.
- . 1981. Taxonomische Notizen zur Gattung *Campylopus* XI. Nova Hedwigia 34: 391–395.
- . 1981. Taxonomische Notizen zur Gattung *Campylopus* VIII. Herzogia 5: 517–529.
- . 1982. Grossdisjunktionen von Arealen südamerikanischer und a afrikanischer Campylopus-Arten. Lindbergia 8: 45–52.
- . 1982. New *Campylopus* records from South and Central America II. Bryologist 85: 322, 323.

329-332.

49: 36-40. \_\_\_\_ & \_\_\_\_ . 1946. Pogonatum flexuosum. Bryologist 49: 141–146. Giese, M. & J.- P. Frahm. 1985 [1986]. A revision of Microcampylopus (C. Müll.) Fleisch. Lindbergia 11: 114-124. \_\_\_\_ & \_\_\_\_. 1985 [1986]. A revision of Campylopodium (C. Müll.) Besch. Griffin, D., III. 1984. A comparison of Breutelia subarcuata (C. Muell.) Schimp. in Besch. and B. chrysea (C. Muell.) Jaeg. in Latin America. Bryologist 87:233-237. \_\_\_\_\_. 1984. Breutelia in Brazil with notes on the occurence of the genus in the New World. J. Hattori Bot. Lab. 57: 83-95. Grout. A. J. 1941. The genus Fissidens in Puerto Rico. Bryologist 44:136-144 \_\_\_\_. 1943. Fissidentaceae. N. Amer. Fl. 15: 167-202, pls. 1-10. \_\_\_\_. 1946. Orthotrichaceae. N. Amer. Fl. 15A:1-62, pls. 1-5. Hampe, E. 1852. Musci frondosi portoricenses, quos Schwanecke, hortulanus, collegit et Hampe examinavit. Linnaea 25: 359-363. Hedwig, J. 1801. Species muscorum frondosorum. Leipzig. Hegewald, E. 1978. Critical notes on Holomitrium (Dicranaceae) from the Antilles. Bryologist 81: 524-531. Higgins, H. H. 1877. Notes by a field-naturalist in the western tropics. Liverpool. [Hoookeria higginsiana Besch. ex Higgins, n. sp., on p. 198] Hooker, W. J. 1818-1820. Musci exotici. 2 vols. London. & R. K. Greville. 1824. Sketch of the characters of the species of mosses belonging to the genera Orthotrichum (including Schlotheimia. Micromitrion, and Ulota), Glyphomitrion, and Zygodon. Edinburgh J. Sci. 1:110-133.

\_\_\_\_. 1983. A monograph of *Pilopogon* Brid. Lindbergia 9:99–116.

. 1985. The world-wide range of Campylopus shawii Wils. J. Bryol. 13:

Frye, T. C. & M. E. Duckering. 1946. Pogonatum convolutum. Bryologist

Husnot, T. 1870. Catalogue des cryptogames recueillis aux Antilles françaises

\_\_\_\_ & \_\_\_\_ . 1825. On the genus *Hookeria* of Smith, of the order Musci.

& \_\_\_\_\_\_. 1825. On the genus Calymperes, Swartz, and Syrrhopodon, Schwägrichen, of the order Musci. Edinburgh J. Sci. 3: 218-227.
Horton, D. G. 1979. Encalypta vittiana sp. nov. and E. flowersiana sp. nov. from North America. Bryologist 82: 368-381. [E. f. from Haiti].
\_\_\_\_\_\_. 1983. A revision of the Encalyptaceae (Musci), with particular reference to the North American taxa. Part II. J. Hattori Bot. Lab.

Edinburgh J. Sci. 2: 221-236.

54: 353-532.


en 1868, et essai sur leur distribution géographique dans ces îles. Bull. Soc. Linn. Normandie II, 4: 19-74. [also as reprint pp. 1-60, pl. I].

- . 1876. Catalogue des muscinées récoltées aux Antilles françaises par T. Husnot. II. Mousses. Rev. Bryol. 3: 57–59, 65–68.
- Jennings, O. E. 1917. A contribution to the botany of the Isle of Pines, Cuba, based upon the specimens of plants from that island contained in the herbarium of the Carnegie Museum under date of October, 1916. Ann. Carnegie Mus. 11: 19–290, pls. V–XXVIII. [mosses pp. 39, 40].
- Jovet-Ast, S. 1947. Mousses épiphytes des Antilles françaises récoltées par P. et V. Allorge en 1936. Rev. Bryol. Lechénol. 15: 154–162.
- Judd, W. S. 1980 [1981]. Notes on the moss flora of Hispaniola. Bryologist 83: 555-558.
- Kallunki, J. A. 1980. Cuban plant collections of J. A. Shafer, N. L. Britton and P. Wilson. Brittonia 32: 397-420.
- Koponen, T. 1979. A synopsis of Mniaceae (Bryophyta). I. South and Central American taxa. J. Ḥattori Bot. Lab. 46: 155–161.
- Lefroy, J. H. 1884. The botany of Bermuda. Bull. U. S. Natl. Mus. 25(2): 35–141. [Sphagnum palustre on p. 126].
- León. 1933. Catalogue des mousses de Cuba. Ann. Cryptog. Exot. 6: 165–214.
- Lewinsky, J. 1984 [1985]. Orthotrichum Hedw. in South America I. Introduction and taxonomic revision of taxa with immersed stomata. Lindbergia 10:65–94.
- Lin, S.-H. 1983. A taxonomic revision of Phyllogoniaceae (Bryopsida). Part I. J. Taiwan Mus. 36(2): 37–86.
- . 1984. A taxonomic revision of Phyllogoniaceae (Bryopsida). Part II. J. Taiwan Mus. 37(2): 1-54.
- Lindberg, S. O. 1863. Om ett nytt slägte, *Epipterygium*, bland bladmossorna. Öfv. K. Vet.-Akad. Förh. 19: 599–609.
- . 1865. Uppställning af familjen Funariaceae. Öfv. K. Vet.-Akad. Förh. 21: 589–608.
- Lorentz, P. G. 1864. Pugillus specierum novarum exoticarum. Pp. 155–167. In: Moosstudien. Leipzig.
- Luan, J. 1814. Mosses. *In*: Hortus jamaicensis; or a botanical description of its indigenous plants hitherto known 1: 512-516. Jamaica.
- Manuel, M. G. 1977. The genus *Meteoridium* (C. Müll.) Manuel, stat. nov. (Bryopsida: Meteoriaceae). Lindbergia 4: 45-55.
- . 1977. Studies in Cryphaeaceae IV. New combinations in Schoeno-bryum. Bryologist 80: 522-524.
- \_\_\_\_\_. 1977. A monograph of the genus Zelometeorium Manuel, gen. nov.

- (Bryopsida: Meteoriaceae). J. Hattori Bot. Lab. 43: 107-126.
- Menzel, M. 1985 [1986]. Die Gattung *Pogonatum* P. Beauv. (Polytrichales, Musci) in Lateinamerika 1. Taxonomie und Geographie von *Pogonatum campylocarpum* (C. Muell.) Mitt. und *P. subflexuosum* (Lor.) Broth. Lindbergia 11: 134–140.
- . 1986. The genus *Pogonatum* P. Beauv. (Musc: Polytrichales) in Latin America 2. Taxonomy and geography of the section *Cephalotricum* (C. Muell.) Besch. Lindbergia 12: 43–46.
- Messmer, L. W. & T. C. Frye. 1947. The *Polytrichum juniperinum* group between South America and the United States. Bryologist 50: 259–268, figs. A, B.
- Miller, H. A. & K.W. Russell. 1975. Key to the mosses to Puerto Rico. Florida Sci. 38: 175–182.
- Mitten, W. 1864. On Anisostichum, a proposed new genus of Musci. J. Linn. Soc., Bot. 7: 119, 120.
- \_\_\_\_\_. 1869. Musci austro-americani. J. Linn. Soc., Bot. 12: 1–659.
- . 1876. The Musci and Hepaticae collected by H. N. Moseley, M. A., naturalist to H. M. S. 'Challenge4.' J. Linn. Soc., Bot. 15: 59-73.
- . 1885. Musci. *In:* Hemsley, W. B., Report on the botany of the Bermudas and various other islands of the Atlantic and southern oceans. First part. Report on the Scientific Results of the Voyage of H. M. S. "Challenger", Botany 1(2): 88–92.
- Mohamed, M. A. H. 1979. A taxonomic study of *Bryum billardieri* Schwaegr. and related species. J. Bryol. 10: 401–465.
- Montagne, C. 1838. Centurie de plantes cellulaires exotiques nouvelles. Ann. Sci. Nat. Bot. II, 9: 38-57.
- . 1843. Quatrième centurie de plantes cellulaires exotiques nouvelles. Décades I–VI. Ann. Sci. Nat. Bot. II, 19: 238–266, pls. 8, 9.
- \_\_\_\_\_. 1845. Cryptogamia. *In:* Sagra, R. de la, Historia física, política y natural de la isla de Cuba 9: 64-328.
- . 1857. Huitième centurie de plantes cellulaires nouvelles, tant indigènes qu'exotiques. Décades IV et V. Ann. Sci. Nat. Bot. IV, 7: 134–153.
- Mullen, D. & T. C. Frye. 1947. Middle and South American species of Oligotrichum. Bryologist 50: 67-79.
- Müller, C. 1843. Macromitrium truncatum nov. sp. Linnaea 17: 383, 384.
- . 1843. De muscis frondosis nonnullis novis vel minus cognitis. Linneae 17: 581–601.
- . 1844. Prionodon. Eine neue Laubmoosgattung. Bot. Zeitung (Berlin) 2: 129–131, 145–148, pl. I.
- . 1846. Beiträge zu einer Flora der Aequinoctial-Gegenden der neuen

pusillus from Martinique and Guadeloupe on p. 2001

Welt. Columbische Laubmoose. Linnaea 19: 193-220. [Zygodon




- 50: 279-310. [Claopodium in Jamaica on p. 292].
- Pessin, L. J. 1922. Epiphyllous plants of certain regions in Jamaica. Bull. Torrey Bot. Club 49: 1–14, pl. I.
- Pursell, R. A. 1960. Fissidens flavosetus, a new species from Haiti. Bryologist 63: 98-101.
- . 1966. Notes on American Erpodiaceae, with descriptions of two new species. Bryologist 69: 461–471.
- . 1971. Nomenclatural notes on American mosses. Bryologist 74: 371–373.
- . 1979. On *Fissidens*. Nomenclatural and distributional notes on some American species. Bryologist 82: 58–64.
- \_\_\_\_\_. 1985. Fissidens buckii and F. quisqueyanus, spp. now. (Bryopsida: Fissidentaceae), from Puerto Rico and the Dominican Republic. Bryologist 88: 74–76.
- . 1986. Typification of Hedwig's species of *Fissidens*. Bryologist 89: 35--41.
- & W. D. Reese. 1963. Studies in American *Fissidens* I. A revision of *F. garberi* Lesq. & James and *F. leptopodus* Card. Bryologist 66: 139–143.
- Questel, A. 1941. The flora of St. Bartholomew (French West Indies) and its origin. [Mosses p. 70 det. E. B. Bartram.]
- . 1951. La flore de la Guadeloupe (Antilles françaises). Paris. ["Les bryophytes ou muscinées" pp. 228–241].
- Reese, W. D. 1959. *Teniola*, a new genus of mosses. Bryologist 62: 221–224.
- . 1961. The genus *Calymperes* in the Americas. Bryologist 64: 89–140. . 1962. *Teniolophora*, a new name for the genus *Teniola*. Bryologist
- . 1962. Teniolophora, a new name for the genus Teniola. Bryologist 65:67.
- . 1977. The genus *Syrrhopodon* in the Americas I. The elimbate species. Bryologist 80: 2–31.
- \_\_\_\_\_. 1978. The genus Syrrhopodon in the Americas II. The limbate species. Bryologist 81: 189–225.
- . 1979 [1980]. New records of Calymperaceae in the Americas. Lindbergia 5: 96–98.
- \_\_\_\_\_. 1983. American *Calymperes* and *Syrrhopodon:* identification key and summary of recent nomenclatural changes. Bryologist 86: 23–30.
- & S. Orbán. 1986 [1987]. Syrrhopodon incompletus Schwaegr. var. incompletus (Musci: Calymperaceae) disjunct between the Americas and Africa. Bryologist 89: 213, 214.
- Reimers, H. 1937. Die europäischen Haplocladium-Arten mit besonderer

Moscosoa 5, 1989 335

Berücksichtigung ihrer aussereuropäischen Verbreitung und ihrer Verwandtschaft. Hedwigia 76: 191–298.

- Renauld, F. & J. Cardot. 1890. Musci exotici novi vel minus cogniti, I. Bull. Soc. Roy Bot. Belgique 29(1): 161–186.
- \_\_\_\_ & \_\_\_\_ . 1894 [1895] . Musci exotici novi vel minus cogniti, VI. Bull. Soc. Roy. Bot. Belgique 33(2): 109-137.
- \_\_\_\_ & \_\_\_\_ . 1904 [1905] . Musci exotici novi vel minus cogniti, X. Bull. Soc. Roy. Bot. Belgique 41(1): 7-122.
- Rendle, A. B. 1936. Notes on the flora of the Bermudas. J. Bot. 74: 101–112 [mosses up. 101, 102, det. H. N. Dixon].
- Rohrer, J. R. 1985. A generic revision of the Hylocomiaceae. J. Hattori Bot. Lab. 59: 241–278.
- Russell, K. W. & H. A. Miller. 1977. The ecology of an elfin forest in Puerto Rico, 17. Epiphytic mossy vegetation of Pico del Oeste. J. Arnold Arbor 58: 1–24.
- Salmon, E. S. 1904. A revision of some species of *Ectropothecium*. Bull. Torrey Bot. Club 31: 309–324, pls. 13, 14 [All species now in *Vesicularia*]:
- Schornherst, R. O. 1943. Phytogeography studies of the mosses of northern Florida. Amer. Midl. Naturalist 29: 509-532.
- Schubert, R. 1978. Beitrag zur Moosflora Kubas. Ergebnisse der ersten kubanisch-deutschen Alexander von Humboldt-Gedächtnis-Expedition 1967/68 der Akademie der Wissenschaften der DDR und Kubanischen Akademie der Wissenschaften Nr. 32. Feddes Repert. 89: 307–326.
- Schwaegrichen, F. B11-1842. Species muscorum frondosorum. Supplementa.. 4 vols. in 11 parts. Leipzig.
- Sharp, A. J., R. Gardner & R. M. Giesy. 1978 [1979]. SEM separation of sterile specimens of *Hedwigidium integrifolium*, *Braunia secunda* and *B. squarrulosa*. Phytomorphology 28: 329–331.
- Shaw, A. J. 1982. *Pohlia* Hedw. (Musci) in North and Central America and the West Indies. Contr. Univ. Mich. Herb. 15: 219–295.
- . 1984. Quantitative taxonomic study of morphology in *Epipterygium*. Bryologist 87: 132–142.
- Sloane, H. 1696. Catalogus plantarum quae in insula Jamaica proveniunt. . . sue prodromi historiae naturalis Jamaicae. Pars prima. London.
- . 1707. A voyage to the islands Madera, Barbados, Nieves, S. Christophers, and Jamaica with the natural history. . . of the last of those islands 1: i-cliv, 1-264, pls. 1-156. London.
- Smith, G. L. 1971. A conspectus of the genera of Polytrichaceae. Mem. New York Bot. Gard. 21(3): 1-83.

- Snider, J. A. 1975. A revision of the genus *Archidium* (Musci). J. Hattori Bot. Lab. 39: 105-201.
- Steere, W. C. 1945. Luisierella, a genus of mosses new to North America. Bryologist 48: 83–85. [Also reported from Puerto Rico].
- . 1946. Mosses of British Honduras and the Department of Petén, Guatemala. III. Bryologist 49: 72–84. [Glossadelphus truncatulus reported from Puerto Rico].
- . 1970. Bryophyte studies on the irradiated and control sites in the rain forest at El Verde. *In*: Odum, H. T. (ed.), A tropical rain forest: a study of irradiation and ecology at El Verde, Puerto Rico. Div. Tech. Info., U. S. Atomic Energy Comm. Pp. D-213-D-225.
- . 1979. The chromosome number of *Callicostella depressa*. Bryologist 82: 103, 104.
- \_\_\_\_\_. 1983. On the continental affiliations of the moss flora of Hispaniola. The World Conference of Bryology, Tokyo, 22–28 May, 1983. Abstracts, p. 28.
- . 1984. The continental affiliations of the moss flora of Hispaniola. J. Hattori Bot. Lab. 56: 19, 20.
- . 1985. On the continental affiliations of the moss flora of Hispaniola. Monogr. Syst. Bot. Missouri Bot. Gard. 11: 155–173.
- Stehlé, H. 1943. La végétation muscinale des Antilles françaises et son intèrêt dans la valorisation sylvicole. Caribbean Forester 4(4). 164–182; 5(1): 20–43.
- Sullivant, W. S. 1861. Musci Cubenses, or mosses collected by Charles Wright in the eastern parts of Cuba during the years 1856, 1857, and 1858. Proc. Amer. Acad. 5: 273–290.
- Swartz, O. 1788. Nova genera & species plantarum seu prodromus descriptionum vegetabilium maximum parte incognitorum que sub itinere in Indiam occidentalem annis 1783–87 digessit. Sweder, Uppsala & Åbo, Stockholm.
- . 1791. Obsevationes botanicae quibus plantae Indiae occidentalis illustrantur emendatur. Erlangen.
- . 1806. Flora Indiae occidentalis aucta atque illustrata sive descriptiones plantarum in prodroko recensitarum 3: 1231–2018. Erlangen.
- Syed, H. 1973. A taxonomic study of *Bryum* capillare Hedw. and related species. J. Bryol. 7: 265–326.
- Taylor, T. 1847. Diagnostic characters of five new species of cryptogamic plants from Jamaica. Ann. Mag. Nat. Hist. I, 20: 379–381. Reprinted in Trans. Bot. Soc. Edinburgh 3: 23–25. 1850.
- Thériot, I. 1916. Une rectification. Bryologist 19: 10.



Wiersma, P. 1984. Moss flora and vegetation of Saba and St. Eustatius (West Indies). Proc. Koninkl. Nederl. Akad. Wetenschappen C, 87: 337-364. Wikstroöm, J. E. 1825. Öfversigt af ön Sanct Barthelemi's flora. Kongl. Svenska Vetenskapsakad. Handl. 1825: 411-433. \_\_\_\_. 1827. Öfversigt af ön Guadeloupe's flora. Kongl. Svenska Vetenskapsakad. Handl. 1827: 51-79. Williams, R. S. 1908. Mosses from tropical America. Bull. Torrey Bot. Club 34: 569-574 ["Cuban mosses collected by W. R. Maxon" pp. 572-574]. \_\_\_\_. 1912. The genus Clastobryum Dozy & Molk, in America. Bryologist 15: 31. \_\_\_\_\_. 1913. Dicranaceae. N. Amer. Fl. 15: 77-158. \_\_\_\_\_. 1913. Leucobryaceae. N. Amer. Fl. 15: 159–166. \_\_\_\_\_. 1919. Archidium cubense sp. nov. Bryologist 22: 2. . 1920. Calymperaceae of North America. Bull. Torrey Bot. Club 47: 367-396, pls. 15-17. . 1921. Hyophila subcucullata sp. nov. Bryologist 24: 22, 23. \_\_\_\_\_. 1921. Mosses from British Guiana and Dominica, Lesser Antilles, collected by Miss E. F. Noel in 1914. Bryologist 24: 65-67. \_\_\_\_\_. 1923. Pilotrichum leoni, sp. nov. Bryologist 26: 50, 51. \_\_\_\_\_. 1927. An undescribed *Taxithelium* from Porto Rico. Bryologist 30: 37 - 39. \_\_\_\_\_. 1929. Macromitrium portoricense sp. nov. Bryologist 32: 69, 70. . 1930. Haitian mosses collected by E. C. Leonard, J. Wash. Acad. Sci. 20: 173-180. Wilson, P. 1917. The vegetation of Vieques Island. Bull. New York Bot. Gard. 8: 379-410 [mosses on p. 409, det. E. G. Britton]. Wilson, W.. 1847. Description of two new mosses from Jamaica. Ann. Mag. Nat. Hist. I, 20: 378, 379. Reprinted in Trans. Edinburgh Bot. Soc. 3: 21, 22. 1850. Zander, R. H. 1977. The tribe Pleuroweisieae (Pottiaceae, Musci) in Middle America. Bryologist 80: 233-269. \_\_\_\_. 1978. Synopsis of the genus *Tuerckheimia* (Pottiaceae); Misc. Bryol. Lichenol. 8: 25-28. \_\_\_\_\_. 1978. A synopsis of Bryoerythrophyllum and Morinia (Pottiaceae) in the New World. Bryologist 81: 539-560. \_\_\_\_\_. 1978. A propaguliferous variant of Oxystegus tenuirostris in Europe, India, Mexico, Cuba and Brazil. Lindbergia 4: 285-288. . 1979. Notes on Barbula and Pseudocrossidium (Bryopsida) in North America and an annotated key to the taxa. Phytologia 44: 177-214. . 1983. A reevaluation of *Neohyophila* Crum (Pottiaceae). Bryologist

Moscosoa 5, 1989 339

86: 134-139.

— & P. M. Eckel. 1982. Hymenostylium recurvirostrum var. insigne and Barbula amplexifolia in British Columbia, Canada. Canad. J. Bot. 60: 1596–1600. [Hymenostylium glaucum var. cylindricum Bartr. from Jamaica transferred to H. recurvirostrum var. cylindricum].

- Zanten, B. O. van. 1959. Trachypodaceae. A critical revision. Blumea 9: 477-575.
- Zündorf, H.-J. 1986. Die Laubmoose Cubas, eine pflanzengeographische Analyse. Abstr. Meeting Centr. Eastern Europ. Bryologists 5: 8, 9.